

- 2011 [Citado 22 Ago 2016]; 12(4): 218-224. Disponible en: <https://www.ncbi.nlm.nih.gov/pubmed/22117753>
29. Nürnberg D, Saboonchi F, Sackey P, Bjorling G. A preliminary validation of the Swedish version of the critical-care pain observation tool in adults. *Acta Anaesthesiol Scand* [Internet] 2011 [Citado 22 Ago 2016]; 55(4): 375 – 506. Disponible en: <http://onlinelibrary.wiley.com/doi/10.1111/j.1399-6576.2010.02376.x/full>
 30. Marmo L, Fowler S. Pain assessment tool in the critically ill post-open heart surgery patient population. *Pain Manag Nurs* [Internet] 2010 [Citado 22 Ago 2016], 11(3): 134-140 Disponible en: <https://www.ncbi.nlm.nih.gov/pubmed/20728062>
 31. Chanques G, Pohlman A, Kress J, Molinari N, Jong A, Jaber S, et al. Psychometric comparison of three behavioural scales for the assessment of pain in critically ill patients unable to self-report. *Crit Care* [Internet] 2014 [Citado 22 Ago 2016]; 18(5): 1-12. Disponible en: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4220092/pdf/13054_2014_Article_2900.pdf
 32. Rijkenberg S, Stilma W, Endeman H, Bosman RJ, Oudemans HM. Pain measurement in mechanically ventilated critically ill patients: Behavioral Pain Scale versus Critical – Care Pain Observation Tool. *J Crit Care* [Internet] 2015 [Citado 22 Ago 2016]; 30(1): 167-172. Disponible en: <http://ezproxy.ufro.cl:2052/science/article/pii/S0883944114003852>
 33. Ministerio de Salud de Chile. MINSAL. Circular Ministerial 2ª/46: Evaluación del Dolor como Quinto Signo Vital. 2004.