

29. Nürnberg D, Saboonchi F, Sackey P, Bjorling G. A preliminary validation of the Swedish version of the critical-care pain observation tool in adults. *Acta Anaesthesiol Scand* [Internet] 2011 [cited en Ago de 2016]; 55(4): 375 – 506. Available from: <http://onlinelibrary.wiley.com/doi/10.1111/j.1399-6576.2010.02376.x/full>
30. Marmo L, Fowler S. Pain assessment tool in the critically ill post-open heart surgery patient population. *Pain Manag Nurs* [Internet] 2010 [cited 22 de Ago de 2016], 11(3): 134-140 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/20728062>
31. Chanques G, Pohlman A, Kress J, Molinari N, Jong A, Jaber S, et al. Psychometric comparison of three behavioural scales for the assessment of pain in critically ill patients unable to self-report. *Crit Care* [Internet] 2014 [cited Ago 2016]; 18(5): 1-12. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4220092/pdf/13054_2014_Article_2900.pdf
32. Rijkenberg S, Stilma W, Endeman H, Bosman RJ, Oudemans HM. Pain measurement in mechanically ventilated critically ill patients: Behavioral Pain Scale versus Critical – Care Pain Observation Tool. *J Crit Care* [Internet] 2015 [cited Ago 2016]; 30(1): 167-172. Available from: <http://ezproxy.ufro.cl:2052/science/article/pii/S0883944114003852>
33. (Ministerio de Salud de Chile. MINSAL. Circular Ministerial 2ª/46: Evaluación del Dolor como Quinto Signo Vital. 2004